The background of the slide features a pulsar, a highly magnetized, rotating neutron star, emitting beams of electromagnetic radiation. The pulsar is depicted with a central bright spot and a surrounding glow of pink and orange light. A white grid of latitude and longitude lines is overlaid on the pulsar image, creating a spherical coordinate system. The entire scene is set against a light blue background.


AGILE & Fermi Observations of PSR B1509-58

Maura Pilia
Università dell'Insubria, Como
INAF-Osservatorio Astronomico di Cagliari


**on behalf of the
AGILE Team & AGILE Pulsar Working Group**

γ incoming gamma ray


Fermi



EGRET




>0.5 GeV Fermi sensitivity
much better than AGILE.
at 100 MeV (and below)
AGILE sensitivity is
competitive.



AGILE

AGILE


20 MeV - 50 GeV



In the 30-100 MeV AGILE sensitivity is competitive (500 cm² eff. Area for timing).

Fermi


20 MeV - 300 GeV



The 1 GeV Fermi sensitivity is much better than the AGILE one.

AGILE Pulsars... three years after...

AGILE detected about 20 gamma-ray pulsars
and tens of candidates from the spatial analysis




AGILE data on 12 Pulsars published so far including
>40% of AGILE Team pulsar targets (AO1 & AO2)

AGILE Pulsars... two years after...

"Discovery of New Gamma-ray Pulsars with AGILE"

(Pellizzoni et al., ApJ, 695, L115, 2009)

July 2007 - June 2008




Many previously unidentified EGRET sources and new AGILE sources are Pulsars!

AGILE Pulsars... two years after...



"Discovery of New Gamma-ray Pulsars with AGILE"

(Pellizzoni et al., ApJ, 695, L115, 2009)

July 2007 - June 2008



Many previously unidentified EGRET sources and new AGILE sources are Pulsars!



$$P \approx 150 \text{ ms}$$


$$\dot{P} \approx 1.53 \times 10^{-12} \text{ s s}^{-1}$$

$$\tau_{sd} \approx 1570 \text{ yrs}$$

$$\dot{E} \approx 1.8 \times 10^{37} \text{ erg s}^{-1}$$


$$B \approx 3.1 \times 10^{13} \text{ G}$$

PSR B1509-58 with Comptel



Kuiper et al. 1999

PSR B1509-58 with AGILE (1)



Pellizzoni et al. *ApJ*, 695, L115, 2009

Pilia et al. *ApJ*, 723, 707, 2010

Detection of PSR B1509-58 and its pulsar wind nebula in MSH 15-52 using *Fermi* - LAT

5





FIG. 2.— Light curves of the pulsar PSR B1509-58 in different energy bands within a circular region of energy-dependent radius. From bottom to the top: COMPTEL (0.75–30 MeV; [Kuitert et al. 1999](#)) and LAT profiles in 30 MeV–300 GeV, 30 MeV–100 MeV, 100 MeV–300 MeV, 300 MeV–1 GeV, 1 GeV–300 GeV energy bands are presented. Two cycles are shown.


Abdo et al., *ApJ*, 714, 927, 2010

PSR B1509-58 with AGILE (2)




The "soft" γ -pulsar B1509-58

Detection of PSR B1509-58 and its pulsar wind nebula in MSH 15-52 using *Fermi* - LAT



AGILE (Pilia et al. 2010)



Fermi (Abdo et al. 2010)

light curves of the pulsar PSR B1509-58 in different energy bands within a circular region of energy-dependent...

Polar Cap - Photon Splitting

Harding, Baring & Gonthier 1997

- Third order QED process
- Forbidden in vacuum
- For $B \leq B_{cr}$ ($B_{B1509} = 3.1 \times 10^{13} G$) it takes place BEFORE the threshold for pair production
- $\gamma \rightarrow \gamma \gamma$ so that high energy emission (> some GeVs) is inhibited

Modelling the Soft Spectrum of PSR B1509-58




FIG. 12.—Same as Fig. 11, for the model spectra of Fig. 8




FIG. 14.—Same as Fig. 13, for the model spectra of Fig. 10

Photon Splitting - Pair Production


$$F(E) \propto E^{-\alpha} \exp[-(E/E_c)]$$
$$\alpha = 1.87 \pm 0.09$$

$$\epsilon_{esc}^{sat} \simeq 0.077 (B' \sin \theta_{kB}, 0)^{-6/5}$$

$$B' = B/B_{cr}$$

θ_{kB} = angle between the photon momentum
and the magnetic field vectors



- Given the observed cutoff energy ($E_c = 81 \pm 20 \text{ MeV}$),
the corresponding magnetic field is $B' = 0.3$. This implies
emission at height $h \simeq 1.3 R_{NS}$ above the PC, where the pair
production ensues.



Pilia et al. 2010

Alternative scenarios

OUTER GAP



Zhang & Cheng 2000

$$\alpha \approx 60^\circ$$

$$\zeta \approx 75^\circ$$

TWO POLE CAUSTIC GAP




FIG. 9. — Collection of sample light curves for the TPC model. Four select w (values in bottom right panel) are shown for each panel; the radio pole has closest approach at phase=0. The values for the number of all peaks, the number of broad peaks and the maximum peak separation (in %) are indicated by each curve. Intensities are normalized to pulse maximum.

Watters et al. 2009




FIG. 14. — Light curves for the Two Pole Caustic (TPC) model. Each panel shows curves for four values of the gap width w . The curves are labeled with the number of major peaks and the peak separation, in percent.

Romani & Watters 2010

$$\alpha \approx 35^\circ$$

$$\zeta \approx 90^\circ$$

Geometry Constraints

OUTER GAP

$$\alpha \approx 60^\circ$$

$$\zeta \approx 75^\circ$$

TWO POLE CAUSTIC GAP

$$\alpha \approx 35^\circ$$



$$\zeta \approx 90^\circ$$

α = magnetic inclination

ζ = viewing angle

- $\alpha < 60^\circ$ at the 3σ confidence level (Crawford et al. 2001)
- If $\zeta > 70^\circ$ (Melatos 1997) then $\alpha > 30^\circ$ at the 3σ level
- For these values, however, the Melatos model for the spin down of an oblique rotator predicts a braking index $n > 2.86$ slightly inconsistent with the observed value $n = 2.839(3)$ (Livingstone et al. 2005)


A Similar Case: PSR B0656+14




$$E_c = 700 \text{ MeV}$$

Weltevrede et al. 2010


New Results from Fermi




$E < 150$
MeV



$E < 300$
MeV



$E < 500$
MeV



Summary

- *AGILE* firmly confirmed the detection of gamma-ray pulsation from PSR B1509-58 with good significance (5 sigma)
- The observed lightcurve shows two peaks which lag the radio peak of 0.39(2) and 0.94(3).
The profile is single peaked up to energies $E = 10$ MeV, where *Comptel* detected the presence of a second peak.
- The detection of pulsed emission by *AGILE* at $E > 30$ MeV, confirms the presence of a strong spectral break, with a cutoff at $E \sim 100$ MeV.

Conclusions

Our observations are compatible with emission from the polar cap regions powered by photon splitting cascades.

This likely interpretation could represent the first physical measurement ever made related to the QED photon splitting process.

The fact that PC emission at HE appears rare might be explained by the requirement that a number of conditions concur to have low magnetosphere emission, e.g. an aligned geometry and high magnetic fields.

New class of "soft" gamma-ray pulsars?

Thank You!

For information and collaborations, please contact us:

Maura Pilia: mpilia@oa-cagliari.inaf.it

Alberto Pellizzoni: apellizz@oa-cagliari.inaf.it