

National Aeronautics and Space Administration



Fermi

Gamma-ray Space Telescope

www.nasa.gov/fermi

Fermi

Gamma-ray Space Telescope

FSSC Data Analysis Workshop




Lifetime and Exposure

E. Hays

Handling Livetime and Exposure

- **LAT specifics**
 - **Calculating exposure is a two-step process**
 - **Livetime: formally the time when the detector is available to collect data**
 - **For analysis, additionally require periods with good data quality and *without* high background**
 - **Large field of view with varying response: need to know **when** and **where****
 - **Exposure: for practical purposes, the number that converts counts to photon flux**
 - **Apply expected photon detection efficiency and angular resolution**
 - **Depends on event selection**
 - **Depends on direction of the photon relative to the instrument boresight and energy**



Livetime



- **Sum up time spent in the field of view for each position in the celestial sky**
 - **Inputs are the photon and spacecraft files**
 - **Options are pixel size and step size for the instrument angles**
 - **Output is a livetime 'cube'**
 - **Respects time-based selection cuts made with gtmktime (GTIs)**

Exposure

- For a position in the sky fold the time spent in each part of the field of view (from livetime cube) with the detector efficiency for that position



P6_V3 effective area is parameterized in theta and energy

Exposure Binning Caveats

- **Response functions are minimally parameterized in the inclination angle (theta) and energy**
 - **Bin or step sizes used in exposure calculations should consider how quickly the efficiency changes in space or energy**
 - **At least 10 bins per decade in energy to avoid errors in estimation where the effective area changes rapidly**
 - **No single, strict recommendation on spatial binning**
 - **1 deg default sufficient for making maps**
 - **Binned likelihood matched to data binning - see threads**
 - **gtexposure invokes a spectral assumption - this has a noticeable impact**
 - **Both the efficiency and angular reconstruction depend on energy**

Exposure Applications

- Each type of analysis has a dedicated method to provide the exposure correction in the appropriate form



- Aperture lightcurve -> `gtexposure`

xxx cm² s

- Unbinned likelihood -> `gtexpmap`

- Binned likelihood -> `gtsrcmaps/gtexpcube`

- All-sky exposure maps -> `gtexpcube`




- See the analysis threads and workshop presentations for details of these applications

Livetime and maximum zenith angle

`gtlucube` knows about the field of view and the spacecraft pointing and can make exposure corrections related to those coordinate systems

if you use
`gtselect zmax=105 ...`
and
`gtmktime ... ROIcut=no`

then you must use
`gtlucube zmax=105 ...`



- If you are doing an all-sky analysis or non-standard zenith cut, the `gtlucube` can make a correction for exposure loss from a zenith angle cut in `gtselect`
 - Note this assumes perfect reconstruction
- If you are doing a basic source analysis and using the standard `ROIcut` in `gtmktime`, you don't need to do this.

Exposure and Zenith Selection Menu

gtmktime inputs

- | | | |
|----------|---|---|
| 1 | GTI filter=DATA_QUAL==1 && LAT_CONFIG==1
ROI cut=yes
<i>Recommended for ToOs and ARR</i> | ROI cut |
| 2 | GTI filter=DATA_QUAL==1 && LAT_CONFIG==1
&& ABS(ROCK_ANGLE)<52
ROI cut=no <i>Used in LAT catalog analysis</i> | Rocking angle cut
(Requires gtlcube
zmax=105) |
| 3 | GTI filter=DATA_QUAL==1 && LAT_CONFIG==1
&& ABS(ROCK_ANGLE)<52
 angsep(RoIRa, RoIDec, RA_ZENITH, DEC_ZENITH)
+ROI_radius<105)
ROI cut=no <i>Least recommended</i> | Rocking angle cut
or ROI cut
(Requires gtlcube
zmax=105) |
| 4 | GTI filter=DATA_QUAL==1 && LAT_CONFIG==1
&& ABS(ROCK_ANGLE)<52
ROI cut=yes

<i>Most conservative and most commonly used</i> | Rocking angle cut
and ROI cut |