



Fermi  
Gamma-ray Space Telescope

# Time evolution of the spectral break in the high-energy extra component of GRB 090926A

***M. Yassine, F. Piron***

*(Laboratoire Univers et Particules de Montpellier - CNRS/IN2P3)*

***F. Daigne, R. Mochkovitch***

*(Institut d'Astrophysique de Paris - CNRS/INSU)*


*On behalf of the Fermi GBM and  
LAT collaborations*

# Outline

---

- **Physical motivations**
- **New time resolved spectral analysis**
  - Best fit model
  - High energy cutoff : significance and temporal evolution
- **Interpretation and new constraints on the jet Lorentz factor**
- **Summary**

# GRB 090926A prompt emission (1/2)



Correlated variability in various bands with a sharp spike at  $T_0 + 10$  s


All energy ranges synchronized (<50 ms)

**Favors internal origin for prompt emission from 10 keV up to GeV energies**


# GRB 090926A prompt emission (2/2)

The emergence of the high-energy power law component coincides with the sharp spike (time bin c)

Cutoff at  $\sim 400$  MeV in time bin c  
If attributed to  $\gamma\gamma$  absorption and used to constrain the jet Lorentz factor :  $\Gamma \sim 200 - 700$



# Luminosity temporal decay



Ackermann 2013 (LAT catalog)

- A break in the temporal decay of the gamma ray luminosity is observed at  $t_b \sim T_0 + 40$  s (observer frame)
- Break time  $t_b$  well after the end of the prompt MeV emission ( $T_0 + 22$  s)
- Afterglow emission after  $t_b$  vs. prompt emission dominated phase before  $t_b$


This work :

- Reanalyse the first 22 s of GRB 090926A with the best data set (Pass 8)
- **Interpretation : Inverse Compton emission /  $\gamma\gamma$  absorption**


# GRB 090926A event statistics at high energies

LAT T90 [5 s-209.8 s]	Pass 7	Pass 8	Pass 8/Pass 7
Number of events	447	1088	2.4
[30 MeV-50 MeV]	33	243	7.4
[50 MeV-100 MeV]	95	381	4.0
[100 MeV-0.5 GeV]	257	391	1.5
[0.5 GeV-1 GeV]	29	40	1.4
[1 GeV-10 GeV]	32	32	1
> 10 GeV	1	1	1


Pass 7



Pass 8



# Best fit model (1/2)




An extra high-energy power law (CUTPL) extending down to  $\sim 10$  keV is not physically motivated (not expected from an IC component)

➔ Use a power law with a break at low energy (Band + CUTBPL)

## Time bin c

Parameters	Band+CUTBPL
<b>Band</b>	
$\alpha$	$-0.94 - 0.02 + 0.03$
$\beta$	$-3.20 - 0.89 + 0.24$
Epeak (keV)	$190 - 9 + 9$
<b>CUTBPL</b>	
Photon index	$-1.48 - 0.08 + 0.09$
E folding (MeV)	$335 - 45 + 65$
C-stat / DOF	$604.7 / 518$
$\Delta$ C-stat w.r.t Band+CUTPL	15




# Best fit model (2/2)

Use a power law with a break at low energy (Band + CUTBPL)

## Time bin d

Parameters	Band+CUTBPL
<b>Band</b>	
$\alpha$	-0.86 - 0.03 + 0.01
$\beta$	-3.1 - 0.5 + 0.2
E <sub>peak</sub> (keV)	177 - 3 + 7
<b>CUTBPL</b>	
Photon index	-1.71 - 0.05 + 0.05
E folding (GeV)	1.20 - 0.18 + 0.22
C-stat / DOF	652.7 / 518
$\Delta$ C-stat w.r.t Band+CUTPL	12



The Band + CUTBPL model fits well the data in the time bins c and d



# Cutoff significance and temporal evolution

Time bins	c [9.8 s, 10.5 s]	d [10.5 s, 21.6 s]	d1 [10.5 s, 12.9 s]	d2 [12.9 s, 21.6 s]
Efolding (MeV)	335 <sup>-45</sup> <sup>+65</sup>	(1.20 <sup>-0.18</sup> <sup>+0.22</sup> ) x10 <sup>3</sup>	550 <sup>-100</sup> <sup>+130</sup>	(1.44 <sup>-0.25</sup> <sup>+0.49</sup> ) x10 <sup>3</sup>
Significance (nb. sigma)	7.6	6.1	4.3	5.1

## Significance of the cutoff

Time bin c :

better constrained

Time bin d :

new spectral break is detected at 1.2 GeV


## Temporal evolution

Time bin c :


no evolution was found

Time bin d :

Increase from 550 MeV (d1) to 1.4 GeV (d2)



# Interpretation 1 : HE break = IC curvature




Comparison to the predictions obtained in the framework of the internal shock model  
*Bošnjak & Daigne 2014*

- MeV component (Band): fast cooling synchrotron radiation, modified by IC in KN regime :  $\alpha_{\text{Band}} \rightarrow -1$
- High-energy component (CUTBPL) :
  - IC with a high-energy shape affected by KN +  $\gamma\gamma$  attenuation
  - Low-energy slope :  $-2 \rightarrow -1.5$

- Example of a single pulse synthetic burst (not adjusted to reproduce GRB 090926A)
- Observed spectral evolution,  $E_{\text{break}}(\text{CUTBPL}) \nearrow$  : KN  $\rightarrow$  Thomson when  $E_{\text{peak}}(\text{Band}) \searrow$  ?
- The comparison with the observed slopes is promising
- The detailed shape (peaks, fluence ratio) is not reproduced yet : a better comparison needs a dedicated simulation of GRB 090926A (ongoing work)

# Interpretation 2 : HE break = $\gamma\gamma$ attenuation

**PRELIMINARY**




- Following *Hascoët et al. 2012*
- $$\Gamma = \Gamma_0(E_{\text{cutoff}}, \Delta t_{\text{var}}) \left[ \frac{1}{2} \left( 1 + \frac{R_{\text{GeV}}}{R_{\text{MeV}}} \right) \left( \frac{R_{\text{GeV}}}{R_{\text{MeV}}} \right) \right]^{-1/2}$$
- $150 < \Gamma_0 < 200$  (similar to Ackermann 2011)
  - $\Gamma$  decreases with increasing ratio between the GeV and MeV emission radii
  - Similar  $\Gamma$  values in the 4 time bins

Time bins (duration)	c (0.7s)	d (11.1s)	d1 (2.4s)	d2 (8.7s)
Ecutoff (MeV)	335	$1.2 \times 10^3$	550	$1.4 \times 10^3$
$\Delta t_{\text{var}}$ (s)	0.15	1	0.5	1

# Summary

---

- GRB 090926A prompt emission has been reanalysed using LAT Pass 8 data
- The attenuation of the high-energy extra PL is detected from  $T_0 + 10$  s to  $T_0 + 22$  s
- With a significant temporal evolution of the break energy from  $\sim 330$  MeV to  $\sim 1.4$  GeV
- Interpretation in terms of SSC internal shock emission is ongoing
- If  $E_{\text{break}}$  attributed to  $\gamma\gamma$  attenuation  $\rightarrow$  new constraints on the jet velocity :  
 $\Gamma \sim 200$  (for  $R_{\text{GeV}}/R_{\text{MeV}}=1$ ) from  $T_0 + 10$  s to the end of the MeV prompt emission



Backup slides


# Black-body component ?

- Adding a BB component for bins c and d does not improve the fit significantly

Preferred model	Parameters	Castor C-stat
<b>Band function</b>		<b>Band</b>
A (ph. cm <sup>-2</sup> . s <sup>-1</sup> . keV <sup>-1</sup> )	0.31 +/- 0.0044	892.07/521
Epeak	278.7 +/- 3.50	
alpha	-0.55 +/- 0.011	
beta	-2.33 +/- 0.0098	
<b>Power law</b>		<b>Band + PL</b>
B (1e-4 cm <sup>-2</sup> s <sup>-1</sup> keV <sup>-1</sup> )	0.38 +/- 0.167	884.61/519
Index	-1.89 +/- 0.088	
Epiv (MeV)	1	
<b>Black-body</b>		<b>Band + BB</b>
Amplitude (1e-6 ph.s <sup>-1</sup> cm <sup>2</sup> keV)	2.92 +/- 0.563	851.06/519
kt (keV)	92.48 +/- 3.02	

- In time bins c and d the best model remains Band + PL\*HighCutoff
- Fitting time bin b with Band + BB model improves the C-stat value by 40 with rmfit (~ 20 with Autofit) w.r.t a Band model.
- In agreement with Guiriec+15 (BB significant at early times)

$$R_{\text{GeV}}/R_{\text{MeV}}$$



$$\Gamma_{\min} \simeq \frac{[C_1 2^{1+2\beta} \mathcal{I}(\beta)]^{1/2(1-\beta)}}{\left[ \frac{1}{2} \left( 1 + \frac{R_{\text{GeV}}}{R_{\text{MeV}}} \right) \left( \frac{R_{\text{GeV}}}{R_{\text{MeV}}} \right) \right]^{1/2}} (1+z)^{-(1+\beta)/(1-\beta)}$$

$$\times \left\{ \sigma_T \left[ \frac{D_L(z)}{c \Delta t_{\text{var}}} \right]^2 E_c F(E_c) \right\}^{1/2(1-\beta)} \left[ \frac{E_{\max} E_c}{(m_e c^2)^2} \right]^{(\beta+1)/2(\beta-1)}$$
(59)

# Time bin c

